

SEND COMPLETED ENROLMENT FORM TO:

**1-855-966-2223** 

Prescriber certification: I certify that this prescription is an original prescription and this pharmacy is the only receiver. The original will not be reused.



support@celltrionconnect.ca



@ Question? 1-855-966-1648



## **Ulcerative Colitis and Crohn's Disease**

|   | d-party providers ("Program Aormation contained in this docun   |  | o facilitate access to Remsima <sup>T</sup>   | <sup>M</sup> SC.   |
|---|---|--|---|--|
| Name: Date of   | of hirth: YYYY/MM/DD Pro  | eferred language: English  | French Other Dlea   | se snecify:  |
| Address:  |   |  |   | se specify.  |
| Tel. (home):  |   |  |   |  |
| Tel. (other):   |   |  | es, please specify:   |  |
| re. (oner).   |   |  |   |  |
| <b>DIAGNOSIS</b> Adult with moderately to severely  | active Crohn's disease  | Adult with moderately  | y to severely active ulcerati   | ve colitis   |
| Other:  |   | Patient completed inducti  | on with intravenous inflixim  | ab? Yes 🗌 No 🗌   |
| VACCINE AND TUBERCULOSIS (TB) ASSESSMENT  TB test   | OPTIONAL TESTING SER  1. Patient switched from Inflito Remsima™ SC maintena  ☐ Therapeutic Drug Monitoring (trough)  ☐ At baseline, before the switch: ☐ At third SC dose ☐ At week ☐ At dose ☐ Repeat inweeks ☐ Repeat inmonths ☐ ASAP ☐ Antidrug Antibody Testin ☐ ASAP   | VICES (PLEASE ALL TH. iximab IV maintenance ince  Calprotectin Testing  IBDoc® OR  QuantOn Cal®  At baseline:  Repeat in weeks Repeat in months  Grand  PHARMACY  Do you have a preferred per pharmacy name:   |   | ximab IV initiation C maintenance Calprotectin Testing   IBDoc® OR   QuantOn Cal®   At baseline:   At first SC dose:   Repeat inweeks   Repeat inmonths  |
| Tel. (office): Fax (office):  |   | Tel.:  |   |  |
| Email:  |   |  |   |  |
| Lilidii.  |   | 16   | Fax:  |  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  |   | Tel  | Fax:  |  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks followin induction regimen with infliximate induction regimen win | sc a <sup>™</sup> SC 120 mg once every ng completion of an ab IV infusion Please  and compl   | • When switching OR to Remsima™ So after the last inf lete the required information.  PRE-MEDICATION ORDER   | Fax:  I from maintenance therapy C, Remsima™ SC may be accelliximab IV infusion  R Please ✓ desired pre-treatre at clinic (indicate dose/route  | y of IV infliximab<br>Iministered 8 weeks<br>nent medication(s)  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks followir induction regimen with infliximate induction regimen wit | sc a™ SC 120 mg once every ng completion of an ab IV infusion Please  and compl   | • When switching OR to Remsima™ SC after the last inf lete the required information. PRE-MEDICATION ORDER administered prior to infusion   | of from maintenance therapy<br>C, Remsima™ SC may be ac<br>diximab IV infusion<br>R Please  | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks followir induction regimen with infliximate induction regimen wit | a <sup>™</sup> SC 120 mg once every ng completion of an ab IV infusion  Please  and compl   | • When switching to Remsima™ SC after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required Pre-medications  Diphenhydramine   | I from maintenance therapy<br>C, Remsima™ SC may be ac<br>liximab IV infusion<br>R Please  desired pre-treatr<br>n at clinic (indicate dose/route<br>Administrat  | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing and the start date:  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Prequency: Inject every 2 weeks  Prequency: Other:  Refills: Other:  For patients transitioning from IV infliximab to Remsima™ Sinfusion:  | a <sup>™</sup> SC 120 mg once every ng completion of an ab IV infusion  Please  and compl   | • When switching to Remsima™ So after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required Pre-medications  | I from maintenance therapy<br>C, Remsima™ SC may be ac<br>liximab IV infusion<br>R Please  desired pre-treatr<br>n at clinic (indicate dose/route<br>Administrat  | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with infliximate with infliximate starts and induction required  • Maintenance dosing of Remsima induction regimen with infliximate a weeks following induction regimen with infliximate starts and induction regimen with infliximate a weeks following induction regimen with infliximate with infliximate substantial properties.  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with infliximate a weeks following induct | a <sup>™</sup> SC 120 mg once every ng completion of an ab IV infusion  Please  and compl   | • When switching to Remsima™ So after the last inflete the required information.      PRE-MEDICATION ORDER administered prior to infusior     □ No pre-meds required Pre-medications     □ Diphenhydramine (e.g., Benadryl*)   | I from maintenance therapy C, Remsima™ SC may be accepted in the second of the second | of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).<br>cion<br>IV 15-30 min prior<br>to infusion (max 50 mg)   |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with infliximab  Dose: 120 mg S  Frequency: □ Inject every 2 weeks Duration: □ Weeks infliximab  Other: □ Weeks starts 4 weeks following induction regimen with infliximab   | a <sup>™</sup> SC 120 mg once every ng completion of an ab IV infusion  Please  and compl   | • When switching to Remsima™ So after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required Pre-medications  Diphenhydramine (e.g., Benadryl*)  Acetaminophen  | I from maintenance therapy C, Remsima™ SC may be ac diximab IV infusion  R Please ✓ desired pre-treatr at clinic (indicate dose/route  Administrat mgmgPO 15-30 minmgmgNV 15-30 min   | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).<br>tion<br>IV 15-30 min prior<br>to infusion (max 50 mg)<br>in prior to infusion<br>prior to infusion  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing the start date:  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Dose: 120 mg S  Frequency: Inject every 2 weeks  Puration:  For patients transitioning from IV infliximab to Remsima™ S  infusion:  Dose induction required  Drug: Intravenous infliximab  Brand:  | sc  a™ SC 120 mg once every ag completion of an ab IV infusion  Please  and completion of an analysis of | • When switching to Remsima™ So after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusion  No pre-meds required  Pre-medications  □ Diphenhydramine (e.g., Benadryl*)  □ Acetaminophen  □ Hydrocortisone   | I from maintenance therapy C, Remsima™ SC may be ac diximab IV infusion  R Please ✓ desired pre-treatr at clinic (indicate dose/route  Administrat mgmgPO 15-30 minmgmgNV 15-30 min   | of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).<br>cion<br>IV 15-30 min prior<br>to infusion (max 50 mg)   |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with inflixing a weeks starts 4 weeks following induction regimen with infliximab subcutaneous)  Dose: 120 mg S  Frequency: □ Inject every 2 weeks Duration: □ Weeks: □ Weeks Duration: □ Weeks: □ Weeks Starts 4 weeks following induction regimen with infliximab subcutaneous)  Dose: 120 mg S  For patients transitioning from IV infliximab to Remsima™ S  Inflixion: □ Dose induction required  Drug: □ Intravenous infliximab  Brand: □ Date of weight: □ Date of w | a™ SC 120 mg once every ng completion of an ab IV infusion  Please  and compl  C  GC, indicate date of last   | • When switching to Remsima™ So after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required Pre-medications  Diphenhydramine (e.g., Benadryl*)  Acetaminophen  Hydrocortisone  Dimenhydrinate  | I from maintenance therapy C, Remsima™ SC may be accelerated by the second sec | of IV infliximab Iministered 8 weeks  ment medication(s) b.  tion  IV 15-30 min prior to infusion (max 50 mg) in prior to infusion prior to infusion  IV 15-30 min prior   |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing the start date:  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Dose: 120 mg S  Frequency: Inject every 2 weeks  Puration:  For patients transitioning from IV infliximab to Remsima™ S  infusion:  Dose induction required  Drug: Intravenous infliximab  Brand:  | a™ SC 120 mg once every ng completion of an ab IV infusion  Please  and compl  C  GC, indicate date of last   | • When switching to Remsima™ SC after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required  Pre-medications  □ Diphenhydramine (e.g., Benadryl*)  □ Acetaminophen  □ Hydrocortisone  □ Dimenhydrinate (e.g., Gravol*)  □ Methylprednisolone  □ Other (indicate):*   | I from maintenance therapy C, Remsima™ SC may be ac Please ✓ desired pre-treatr n at clinic (indicate dose/route  Administrat  mg PO 15-30 min mg PO or □  mg PO or □  mg PO or □  mg PO or □   | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).<br>Stion<br>IV 15-30 min prior<br>to infusion (max 50 mg)<br>in prior to infusion<br>prior to infusion<br>IV 15-30 min prior<br>to infusion<br>or to infusion  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing the start date:  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Dose: 120 mg S  Frequency: Inject every 2 weeks  Por patients transitioning from IV infliximab to Remsima™ S  Infusion:  Dose induction required  Drug: Intravenous infliximab  Brand:  Patient weight: Date of weight:  Dose (mg): Exact dose: OR Exact # of vials:  Frequency/duration   | a™ SC 120 mg once every ng completion of an ab IV infusion  Please  and compl  C  GC, indicate date of last   | • When switching to Remsima™ SC after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required  Pre-medications  □ Diphenhydramine (e.g., Benadryl*)  □ Acetaminophen  □ Hydrocortisone  □ Dimenhydrinate (e.g., Gravol*)  □ Methylprednisolone  □ Other (indicate):*   | I from maintenance therapy C, Remsima™ SC may be accelerated by the second sec | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).<br>Stion<br>IV 15-30 min prior<br>to infusion (max 50 mg)<br>in prior to infusion<br>prior to infusion<br>IV 15-30 min prior<br>to infusion<br>or to infusion  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing subcutaneous.  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Dose: 120 mg S  Frequency: Inject every 2 weeks Duration:  For patients transitioning from IV infliximab to Remsima™ S  infusion:  Dose induction required  Drug: Intravenous infliximab  Brand:  Patient weight: Date of weight:  Dose (mg): Exact dose: OR Exact # of vials:  Frequency/duration   | SC  a™ SC 120 mg once every g completion of an ab IV infusion  Please  and completion  C  GC, indicate date of last  YYYYY/MM/DD  100 mg vials  | • When switching to Remsima™ SC after the last inflete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required  Pre-medications  □ Diphenhydramine (e.g., Benadryl*)  □ Acetaminophen  □ Hydrocortisone  □ Dimenhydrinate (e.g., Gravol*)  □ Methylprednisolone  □ Other (indicate):*   | I from maintenance therapy C, Remsima™ SC may be ac Please ✓ desired pre-treatr n at clinic (indicate dose/route  Administrat  mg PO 15-30 min mg PO or □  mg PO or □  mg PO or □  mg PO or □   | y of IV infliximab<br>Iministered 8 weeks<br>ment medication(s)<br>).<br>Stion<br>IV 15-30 min prior<br>to infusion (max 50 mg)<br>in prior to infusion<br>prior to infusion<br>IV 15-30 min prior<br>to infusion<br>or to infusion  |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing the start date:  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Dose: 120 mg Start date:  Prequency: Inject every 2 weeks Duration:  For patients transitioning from IV infliximab to Remsima™ Start date:  Pose induction required  Drug: Intravenous infliximab  Brand:  Patient weight: Date of weight:  Dose (mg): Exact dose: OR Exact # of vials:  Frequency/duration  IV Induction weeks: 0   2   6   Other dosing in the start date:   | SC  a™ SC 120 mg once every g completion of an ab IV infusion  Please  and completion of an ab IV infusion  Please  and completion of an analysis of analysis of an analysis  | • When switching to Remsima™ So after the last infete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required Pre-medications  Diphenhydramine (e.g., Benadryl*)  Acetaminophen  Hydrocortisone  Dimenhydrinate (e.g., Gravol*)  Methylprednisolone  Other (indicate):*  "Please provide patient with a prescription to the Program and to secreption to the Program and to secreption to the Program and to secreption to the Program and to | I from maintenance therapy.  C, Remsima™ SC may be accellized by infusion  R Please  desired pre-treatre at clinic (indicate dose/route  Administrate  mg  PO or   | of IV infliximab Iministered 8 weeks Iment medication(s) (b).  Sion  IV 15-30 min prior to infusion (max 50 mg) in prior to infusion  IV 15-30 min prior to infusion  IV 15-30 min prior to infusion  IV 15-30 min prior to infusion  Iverproper to infusion (max 50 mg) in prior (max 50 mg) in pr |
| PHYSICIAN PRESCRIBING SECTION FOR REMSIMA™  DOSAGE AND ADMINISTRATION  • Maintenance dosing of Remsim 2 weeks starts 4 weeks following induction regimen with inflixing.  Requested start date:  Prug: Remsima™ SC (infliximab subcutaneous)  Autoinjector: (DIN: 02511584)  Dose: 120 mg S Prequency:  Inject every 2 weeks Duration:  Refills:  Other:  For patients transitioning from IV infliximab to Remsima™ Sinfusion:  Dose induction required  Drug: Intravenous infliximab  Brand:  Patient weight:  Dose (mg): Exact dose:  OR Exact # of vials:  Frequency/duration  IV Induction weeks:  O  | SC  a™ SC 120 mg once every g completion of an ab IV infusion  Please  and completion of an ab IV infusion  Please  and completion of an analysis of analysis of an analysis  | • When switching to Remsima™ So after the last infete the required information.  PRE-MEDICATION ORDER administered prior to infusior  No pre-meds required Pre-medications  Diphenhydramine (e.g., Benadryl*)  Acetaminophen  Hydrocortisone  Dimenhydrinate (e.g., Gravol*)  Methylprednisolone  Other (indicate):*  "Please provide patient with a prescription to the Program and to secreption to the Program and to secreption to the Program and to secreption to the Program and to | I from maintenance therapy.  C, Remsima™ SC may be accelliximab IV infusion  R Please  desired pre-treatment at clinic (indicate dose/route)  Administration  mg  PO 0 r  PO or  | of IV infliximab Iministered 8 weeks Iment medication(s) (b).  Sion  IV 15-30 min prior to infusion (max 50 mg) in prior to infusion  IV 15-30 min prior to infusion  IV 15-30 min prior to infusion  IV 15-30 min prior to infusion  Iverproper to infusion (max 50 mg) in prior (max 50 mg) in pr |

| NOTES   |   |
|---|---|
|   |   |
|   | _   |
|   |   |
|   |   |
| PATIENT CONSENT   |   |
| The CELLTRION CONNECT™ Patient Support Program (the "Program") is a patient support program been prescribed Remsima™ SC. The Program services may include health/disease/product information or financial assistance (the "Support Services"). A third-party service provider, McKesson Canada Cor Administrator"). Its employees and/or agents handle your personal information, which is processed in protection standards, as may be designated from time to time by Celltrion.                    | on, insurance reimbursement assistance, treatment services rporation, is the administrator of the Program ("Program |
| (1) that personnel of the Program Administrator ("Program Personnel") may contact me by any me  | eans (e.g., phone, fax, email, mail, etc.) for the purposes of  |
| <ul> <li>administering the Support Services;</li> <li>(2) that my personal health information may be collected, used and stored by the Program Administrate delivery of the Support Services;</li> </ul>  | strator and by my healthcare providers involved in  |
| (3) that my personal information may be exchanged among Program Personnel, my healthcare pro<br>or Celltrion's agents and service providers, such as information technology providers, for purpos<br>Support Services; and  |   |
| (4) that my healthcare providers and the Program Administrator may share my personal information its legal and regulatory obligations, including with respect to safety and adverse drug reporting  |   |
| I understand that the Program Administrator may also share de-identified information (i.e., where per (combined with other data) with Celltrion to conduct analyses for commercial, market and scientific or as otherwise may be permitted by law.  |   |
| I understand that the collection, use and disclosure of information contemplated herein may involve to utside of Canada (including in the United States), where local laws may require the disclosure of per circumstances that are different than those that apply in Canada. The reasonable contractual measure processed or handled by these third parties outside of my country of residence may be subject to for to disclose personal information to government authorities in those countries. | rsonal information to governmental authorities under<br>res taken to protect my personal information while          |
| I understand and agree that Celltrion has the right without notice to (1) make changes to the scope of the eligibility requirements for the Support Services; or (3) discontinue the Program or any of the Support Services.  |   |
| If at any time Celltrion appoints a new program administrator, I will be notified of same and I hereby at the new program administrator for the purposes of continuing my participation in the Program.   | authorize Celltrion to transfer my personal information to  |
| I understand that I have the right to have access to or to correct my personal information held by Prolocated at 4705 Dobrin, Saint-Laurent, Quebec, H4R 2P7, and by telephone at: 1-855-966-1648.  |   |
| I understand that I have the right at any time to withdraw my consent to the use of my personal infor participating in the Program.  In signing this form, I consent to the above.  | mation but in 1 do décide to do so, i wiii no longer be   |
| $\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $  | nic or other means for the purposes of market research.   |
| I acknowledge that I may at any time opt-out from such communications by advising Program Ac  | dministrator by email at: support@celltrionconnect.ca   |
|   |   |
| Patient signature:  | YYYY/MM/DD  |





**6** 1-855-966-1648



1-855-966-2223



support@cell trion connect.ca



www.celltrionconnect.ca



Celltrion Healthcare Canada Limited 121 King Street West, Suite 1010 Toronto, Ontario M5H 3T9 www.celltrionhealthcare.ca